## Improving the soil

The ideal vegetable garden soil is deep, friable, well-drained, and has high organic matter content. Proper soil preparation provides the basis for good seed germination and subsequent growth of garden crops. Careful use of various soil amendments can improve garden soil and provide the best possible starting ground for your crops.

## Tilling the soil

It was once assumed that garden soil should be turned over completely every year, mostly for weed and pest control. More recent ndings show that this practice is actually detrimental in some cases, causing soil compaction, upsetting balances of microorganisms, and often causing layers of course organic material to be buried below the in uence of insects and microbes that would otherwise cause breakdown of the material.

Rotary tilling (roto-tilling) is sui cient for most home gardens as long as plant debris accumulation is not out of hand. Rotary tilling mixes the upper layers of soil rather than completely turning the soil over, and the f ects produced are generally desirable. One possible harmful f ect of roto-tilling is the formation of a compaction layer just beyond the reach of the tines. Use of deep-rooted cover crops or double-digging can do much to prevent or alleviate this problem. Of course, small gardens can be worked entirely by hand using a spade or garden fork.

Gardeners often wonder whether to till in the spring or fall. Working the soil in the fall has several advantages. It allows earlier spring planting, since the basic soil preparation is already done when spring arrives. Turning under large amounts of organic matter is likely to result in better decomposition when done in the fall since autumn temperatures are higher than those of early spring, and there is more time for the process to take place. Insects, disease organisms, and perennial weeds may be reduced by killing or inactivating them through burial or exposure to harsh winter weather. The physical condition of heavy clay soils may be improved by the alternate freezing

and thawing, which breaks up tightly aggregated particles. Also, snow is trapped between the hills of ploughed soil, so more moisture is retained than on\_at, bare ground. Incorporation of limestone or rock fertilizers in the fall gives them time to become integrated with the soil and in\_uence spring plant growth.

Fall tilling alone is not recommended for hillside or steep garden plots, since soil is left exposed all winter, subject to erosion when spring rains come. If a winter cover crop is grown to improve soil and prevent erosion, the ground will have to be tilled in the fall to prepare the soil for seed and again in spring to turn under the green manure. Spring cultivation is better for sandy soils and those where shallow tilling is practiced.



Roto-tilling is usually su cient for most gardens, and saves labour. However, be careful not to over-till, as this can destroy soil structure. © Lana Bos, Dalhousie

1

Whatever the method or timing of tillage, it is extremely important to work only when soil moisture conditions are right. To test, pick up a handful of soil and squeeze it. If it stays in a ball, it is too wet. If it crumbles freely, it should be about right. Excessively dry soil is powdery and clumpy and may be di cult to work. If soil sticks to a shovel or if when spading, the turned surface is shiny and smooth, it is still too wet. Working wet soil can destroy structure, which may take years to rebuild. Ploughing with a tractor when the soil is wet is especially damaging, causing the



formation of a compaction layer that will inhibit root growth.

Just prior to planting, break up large clods of soil and rake the bed level. Small-seeded vegetables germinate

best in smooth, ne-surfaced soil. Do not pulverize the seedbed by over tilling. This destroys the structure and promotes crusting and erosion.

Research into no till' gardening systems indicates ing and erosion.



Organic matter is a great soil improver for both clay and sandy soils. Good sources of organic matter include manures, leaf mould, and straw. These materials are decomposed in the soil by soil organisms. Various factors, such as moisture, temperature, and nitrogen availability, determine the rate of decomposition through their  $\mathfrak e$  ects on the organisms. Adequate water must be present, and warm temperatures will increase the rate at which the microbes work. The proper balance of carbon and nitrogen in the material is needed to ensure adequate nutrient availability both to growing plants and the microbes. Adding nitrogen may be necessary if large amounts of undecomposed leaves, straw, or other high-carbon substances are used.

The use of compost is one way to get around the decomposition problem. Compost is usually made by the gardener from plant and/or animal wastes. The basis of the process is the microbial decomposition of mixed, raw, organic materials to a dark, uf y product resembling rich soil, which is worked into the garden soil (see Composting' in this guide).

Animal manures are commonly used as a garden soil amendment. The nutrient content of manure varies. Fresh horse, sheep, rabbit, and poultry manures are quite high in nitrogen and may even burn plants if



High quality compost is decomposed before it is added to the soil.

© Andy
Hammermeister,
Dalhousie.

applied directly to a growing garden. They are best applied in early fall and tilled under. Manure usually has fewer total nutrients than synthetic fertilizers in terms of N, P and K, but is a valuable soil builder. Unfortunately, manures may be a source of weed seeds if this is a problem, composting in a hot pile may help. In urban areas, manure can be purchased as a bagged product sold at garden centers.

Another source of inexpensive soil builder is the cover crop. Green manures, or cover crops, such as annual rye, ryegrass, buckwheat, and oats, are planted in the

garden in the late summer for incorporation in the spring. For best results, seed should be sown four to six weeks before the rst frost. In a fall garden, plant cover crops between the rows and in any cleared areas. Cover cropping provides additional organic matter, holds nutrients that might otherwise be lost over winter, and helps reduce erosion. Legume cover crops can increase the amount of nitrogen in the soil and reduce fertilizer needs. A deep-rooted cover crop allowed to grow for a season in problem soil can help break up hardpan and greatly improve tilth. Incorporate green manures at least two weeks before planting vegetables. They should not be allowed to go to seed before incorporation or you will have a weed problem.

The regular addition of manure, compost, cover crops, and other organic materials can raise the soil

nutrient and physical level to a point where synthetic fertilizers are no longer needed. This happens through the intrinsic fertilizing value of the amendment, but also through the increased action of microorganisms on soil and humus particles humic acid (and other acids) helps to release previously locked-up nutrients present in the soil, and the extra surface area provided by humus serves as a reserve, holding nutrient elements until they are needed by plants. This highly desirable soil quality does not come about with a single or even several additions of organic material but rather requires a serious, long-term soil building program. Soil is the base for plant growth and much attention should be paid to getting and keeping it in the best condition possible.



## Activity

## Familiarize yourself with soil improving products

Use the table to list ve of the soil amendments presented in this factsheet, particularly those that address issues you may have in your garden soil. Note what soil condition(s) the amendment is used for. Look for sources of these amendments gardening stores, catalogues, websites, local farmers, etc. Also note the cost of each amendment.

Amendment	Use	Available from	Cost
1.			
2.			
3.			
4.			
5.			